


# MEDICAL MISSIONARY WORK

AT CESAREA, TURKEY



*Cesarea and Mt. Argeris*

BY ALDEN R. HOOVER, M. D.

FOR YEAR ENDING AUGUST 31

1912

# American Hospital

TALAS, CESAREA, TURKEY IN ASIA

## Hospital Staff

<i>Physician in charge</i>	A. R. HOOVER, M. D.
<i>Assistant Physician</i>	G. K. GULBENK, M. D.
<i>Anæsthetist</i>	MR. HAIG HARATUNIAN, Ph.G.
<i>Superintendent</i>	MISS THEDA B. PHELPS
<i>Head Nurse</i>	MISS MELPOMENE SERINIDOU
<i>Surgical Nurse</i>	MISS YEGHSAPET BJOYAN
<i>Nurse in training</i>	MISS SERANOUSH BOYAJIAN
<i>Nurse in training</i>	MISS LUCIA SARKISIAN
<i>Nurse in training</i>	MISS YESTER BOSHNAKIAN
<i>Housekeeper</i>	MISS EPISTEME YOUAKIMIDOU
<i>Night Nurse</i>	MRS. THEOBISTA ANANIADOU
<i>Night Nurse</i>	MRS. YEGHSAPET SARRAFIAN
<i>Orderly</i>	MR. NAZAR HAJIKIAN
<i>Orderly</i>	MR. LEVON SELVERIAN

## Dispensary

<i>Dispenser</i>	MR. HAIG HARATUNIAN, Ph.G.
<i>Assistant dispenser</i>	MR. YORIYOS YOUAKIMIDOU, B. A.

## Medical Missionary Work

---

**T**ALAS, a town of 10,000 inhabitants is situated practically in the center of Asia Minor. It is one and one half hours from the city of Cesarea, and at an altitude of 4,300 ft. above sea-level. It commands a beautiful view of Mt. Argeus on the one hand, and the broad and fertile Cesarea plain on the other.

Medical work was started in Talas twenty years ago by Dr. William S. Dodd.



TALAS

Through his untiring efforts, combined with those of Dr. Wilfred Post, his associate during the last seven years of this time, the work was built up in a permanent and substantial manner and recognized throughout the country as one of the most important medical centers in Turkey. A splendid three-story stone Hospital building was erected also a two-story stone dispensary block with rooms for drug-store and a commodious suite of five waiting rooms and examination rooms. That these men were wise

in the selection of Talas as a medical center, and that their careful and efficient work was held in high esteem is attested by the following figures showing the increasing clientele the past four years.

	1908-09	1909-10	1910-11 (½ year only)	1911-12
Number out patients . . . . .	2691	3433	1818	3832
Number in-patients . . . . .	419	460	197	616
Number surgical operations . .	667	691	299	785

For some years past the former doctors have contemplated starting a new work in Konia and one and one half years ago the plans were completed whereby the

American Board took over the Hospital and dispensary buildings and grounds together with a part of the former equipment, and the Talas medical work passed entirely into the hands of that organization.



ENTRANCE TO HOSPITAL

These changes necessitated closing the Hospital for a period of six months, an entire change in the personnel of the Hospital staff, and the providing of a new equipment.

Dr. A. R. Hoover, having had five years experience in the American Board's

hospital at Marsovan, was secured as physician in charge. Miss Theda B. Phelps, graduate of Illinois Training School, and afterward superintendent of Colonial Hospital, Geneva, Illinois, was secured as superintendent. Dr. Gulbenk, graduate of Anatolia Hospital, Marsovan, and of Beirut College of Medicine was secured as assistant physician; Mr. Haig Haratunian, graduate of Beirut College of Pharmacy, as anæsthetist and dispenser. These together with one trained Armenian nurse, and a corps of undergraduate nurses, orderlies, servants, and helpers made up the hospital staff and employees.

It seemed that all things were to work together for good for Talas, for although a multitude of difficulties stood in the way of re-establishing the work, one by one

they have been in great part overcome, leaving many, but such as are to be expected in any extensive work in this country. The matter of securing an equipment was a most important one, and this seemed providentially met, for there stood waiting, stored in Constantinople, an outfit of some two thousand to three thousand dollars' value practically unused, which had been purchased for an American Hospital in Constantinople. This outfit through the kindness of Constantinople station was secured and shipped to Talas. The dispensary was restocked with drugs and supplies from London, and after a period of inactivity of six months the Talas Hospital was re-opened Sept. 1, 1911.

The work started out at once in good force, and before many days the hospital was filled as far as its equipment would allow. The heavy task of the fall was succeeded by steady service throughout the winter months, the winter being of rather exceptional mildness. This was followed by a rush of work in the spring which continued until July, when as has been customary the Hospital was closed for a few weeks for repairs and to give a brief respite from strenuous duty during the most heated term. The Hospital was open for patients 325 days throughout the year. The capacity of the Hospital, formerly forty beds, was of necessity increased to seventy beds. In this manner some 150 more in-patients have been accommodated than formerly. This increase together with the small Hospital force has necessitated the omission of much work which should be done. Practically no touring work has been possible; professional visits have been made to some twelve cities and villages of this region. Much needed relief-work for poor has had to be neglected, while many plans for extension of the work in various lines have failed of execution for lack of time. During the year we have been in constant dread of cholera; it is still raging within four hours of Talas, but so far the cities of Talas and Cesarea have escaped. An epidemic of measles last winter carried off some hundreds of young children due to lack of care during the period of convalescence. An incident of unusual occur-



"THE POOR YE HAVE ALWAYS WITH YOU"—IN TURKEY

rence in this country was a mad-dog scare. Some twenty-four children of Talas were bitten. These children were taken in hand by the Government officials and sent to Constantinople where they were given serum treatment for a period of fourteen days. Among the Hospital and out-patient cases have been a number of general medical interest, and these have been reported in current medical literature.

The constituency of the Hospital is the most varied imaginable, from the beggar in filth and rags, to those prominent in civil and religious circles. As to race and religion, about one third are of Moslem faith and two thirds of Christian races while the promiscuous range of work is perhaps best seen by reference to the list of surgical operations appended. The unsettled political atmosphere has not affected work here in the interior to any material degree. Emigration to America from the cities and villages of the whole region continues unabated, affecting particularly the Christian races, Armenian, and Greek. This will mean an increasing proportion of Moslem patients from year to year.



THE DISPENSARY

institution of such high standard of proficiency as shall command the confidence and respect of Moslem and Christian alike. There is an attempt to set an honest standard of medical ethics in the midst of much to the contrary, and first of all to gain confidence, then changed lives. Aside from personal talks with patients, daily morning and evening prayers are held in the Hospital wards, and a short preaching service each Sabbath afternoon.

A training school for nurses will be started again after a lapse of two years. There are three candidates for the beginning class, the course of instruction to be three years. Graduates of schools of similar grade to the girls' school at Talas will be eligible to the training classes, a working knowledge of English being essential.

The distinctively religious aspect of the medical missionary work has been done in a purposely quiet and unassuming manner. It seems wise to us and consistent with the policy of our Board to attempt to build up a medical

The gratitude attested by many is our best evidence of the hand of God in all the work.

We acknowledge with gratitude the faithful and efficient service of our pastor, Rev. Mibar Muncherian, in conducting evening devotions in an exceedingly helpful manner. We would also gratefully acknowledge the service of Mr. Vartavar Garabedian in conducting the Sabbath afternoon service in the men's ward. Services in the women's wards have been conducted by various persons, and to them also our thanks are due, as well also to our missionary fellow-workers of Talas station.

Financially the Hospital has not yet reached the self-supporting ideal, although the returns from patients have been encouraging in this field where the rich are few and the poor abound.

The re-opening of the work this year was naturally attended by numerous incidental expenses; the American Board has generously

made a grant of \$2,000 to help the financial strain. It is sincerely hoped that many friends in America or elsewhere will see in this work at Talas an opportunity for investment which is yielding great returns. A number have signified their desire to contribute \$150 a year for the support of one bed each. Subscriptions which have so far reached us we acknowledge with thanks.



COMING TO THE DISPENSARY

Mrs. Horace Crawford, Winnipeg,	.	.	.	.	.	.	\$100
Mrs. James H. Ashdown, Winnipeg	.	.	.	.	.	.	50
Dr. C. A. Noland, Ogden, Iowa	.	.	.	.	.	.	50





## Surgical Operation List.

<i>Regional Surgery</i>		<i>Gynecological</i>	
Abdominal		Curettage . . . . .	27
Appendectomy . . . . .	16	Curettage and cervical repair . . . . .	5
Gastro-Enterostomy, posterior . . . . .	9	Adherent placenta . . . . .	2
Entero-Enterostomy . . . . .	1	Abortion 5 mos. foetus . . . . .	1
Volvulus intestines . . . . .	2	Cervical repair . . . . .	1
Cholecystotomy . . . . .	2	Perineomaphy . . . . .	7
Ventral Hernia . . . . .	6	Vesico-vaginal fistula . . . . .	5
Stab wound prolapse omentum . . . . .	2	Recto-vaginal fistula . . . . .	1
Stab wound prolapse intestine, gangrene . . . . .	1	Embryotomy . . . . .	2
Appendicitis suppurative . . . . .	6	Forceps delivery . . . . .	1
Tubercular peritonitis . . . . .	11	Pelvic abscess vaginal . . . . .	10
Exploratory laparotomy . . . . .	3	Pelvic abscess groin . . . . .	2
Hysterectomy, complete inversion uterus . . . . .	1	Condylomata vaginæ . . . . .	1
Hysterectomy uterine and ovarian fibroids . . . . .	1		—
Hysterectomy abdominal . . . . .	5	Total gynecological . . . . .	65
Ventral fixation or suspension . . . . .	7		
Ventral fixation and ovariectomy . . . . .	1	<i>Eye, Ear, Nose, and Throat</i>	
Ventral fixation and appendectomy . . . . .	1	Cataract . . . . .	26
Ovariectomy single or double . . . . .	3	Dacryocystitis . . . . .	8
Resection ovary . . . . .	1	Entropion single or double . . . . .	40
Ovariectomy and appendectomy . . . . .	5	Knapp's Operation . . . . .	13
Pyosalpinx . . . . .	2	Enucliation . . . . .	10
Pelvic adhesions . . . . .	1	Iridectomy . . . . .	9
Pelvic adhesions and appendectomy . . . . .	1	Soft cataract . . . . .	4
Resection Fallopian tube . . . . .	1	Dislocation lens . . . . .	3
Dermoid cyst ovary . . . . .	1	Pterygium . . . . .	6
Adherent ovarian cyst . . . . .	1	Ectropion . . . . .	6
	—	Strabismus . . . . .	4
Total abdominal . . . . .	91	Retention cysts . . . . .	5
		Abscess eyelid . . . . .	1
<i>Hernias</i>		Tonsils and adenoids . . . . .	12
Hernia inguinal single or double . . . . .	36	Radical mastoid . . . . .	5
Hernia inguinal female . . . . .	3	Perforation palate . . . . .	5
Hernia inguinal strangulated . . . . .	1	Deviation nasal septum . . . . .	3
Hernia inguinal strangulated gangrenous . . . . .	1	Specific abscess septum . . . . .	1
Hernia inguinal and hydrocele . . . . .	1	Nasal polyp . . . . .	1
Hernia inguinal and castration . . . . .	1	Sarcoma of nose . . . . .	1
Hernia inguinal and hernia bladder . . . . .	1	Laceration ear . . . . .	3
Hernia congenital . . . . .	3	Caries external auditory canal . . . . .	1
Hernia femoral . . . . .	1	Caries middle ear . . . . .	1
Hernia femoral strangulated . . . . .	1	Pharyngeal abscess . . . . .	1
Hernia femoral strangulated gangrenous . . . . .	1		—
Hernia direct . . . . .	1	Total eye, ear, nose, and throat . . . . .	169
Hernia double with traumatic laceration of hernial sac . . . . .	1	<i>Calculi</i>	
	—	Suprapubic cystotomy . . . . .	10
Total hernias . . . . .	52		—
		Total calculi . . . . .	10

Amputations		Ileum . . . . .	1
Fingers . . . . .	9	Ilio-sacral joint . . . . .	2
Hand . . . . .	1		
Forearm . . . . .	1	Total tbc. bones and joints . . . . .	64
Arm . . . . .	2		
Charcot foot . . . . .	1	Tubercular Sinuses and Abscesses	
Symes foot . . . . .	3	Scalp . . . . .	1
Leg . . . . .	5	Cervical glands . . . . .	20
Thigh . . . . .	2	Cervical and axillary . . . . .	8
		Cervical and groin . . . . .	1
Total amputation . . . . .	24	Axillary glands . . . . .	11
		Thigh sinus and abscesses . . . . .	13
Gunshot		Arm . . . . .	2
Face, eye, and hand . . . . .	1	Elbow . . . . .	1
Hand and wrist . . . . .	1	Groin glands . . . . .	1
Arm and forearm . . . . .	8	Leg . . . . .	1
Scapula . . . . .	1	Chest wall . . . . .	2
Back of neck . . . . .	1	Hip . . . . .	1
Pubic bone and bladder . . . . .	1	Popliteal abscess . . . . .	2
Arm, ligation brachial artery . . . . .	1	Foot . . . . .	1
Both femurs . . . . .	1	Loin . . . . .	1
Thigh . . . . .	5		
Knee . . . . .	3	Total tbc. sinuses and abscesses . . . . .	66
Leg and foot . . . . .	3		
Groin . . . . .	2	Tumors	
Total gunshot . . . . .	28	Epithelioma face . . . . .	4
		Lipoma . . . . .	5
Genito-Urinary		Cancer lip with or without glands . . . . .	5
Urinary fistulae . . . . .	4	Cancer breast and axillary glands . . . . .	4
Posterior urethrotomy . . . . .	3	Cancer scalp and neck . . . . .	1
Internal urethrotomy . . . . .	5	Cancer forehead plastic . . . . .	1
Hydrocele double . . . . .	1	Cancer sternum . . . . .	1
Castration single or double . . . . .	4	Cancer perineum . . . . .	1
Hypospadias . . . . .	3	Cancer bladder, supra-pubic . . . . .	1
Plastic scrotum . . . . .	1	Cancer nose, amputation . . . . .	1
Nephrectomy . . . . .	1	Cancer orbit . . . . .	2
Total genito-urinary . . . . .	22	Sarcoma-melano, forehead . . . . .	1
		Sarcoma axilla . . . . .	1
Tubercular Bones and Joints		Sarcoma testes . . . . .	1
Lacrymial bone . . . . .	1	Sarcoma occiput bone . . . . .	1
Superior maxillary . . . . .	3	Sarcoma thigh . . . . .	1
Inferior maxillary . . . . .	1	Thyroidectomy . . . . .	4
Orbit and cranium . . . . .	1	Fibroma rectum . . . . .	1
Ribs, resection . . . . .	5	Epithelioma back . . . . .	1
Ribs, without resection . . . . .	1	Gummata . . . . .	1
Humerus . . . . .	3	Fibroma forearm . . . . .	1
Glenoid cavity . . . . .	1	Epulis upper jaw . . . . .	1
Radio-humeral joint . . . . .	1	Keloid multiple . . . . .	3
Resection elbow joint . . . . .	2	Papilloma multiple foot . . . . .	1
Resection knee joint . . . . .	16	Angioma foot . . . . .	1
Synovitis knee joint . . . . .	4		
Hand and wrist . . . . .	4	Total tumors . . . . .	45
Hand and foot . . . . .	1		
Synovitis ankle . . . . .	3	General Surgery	
Ankle bones . . . . .	7	Lacerated wound scalp . . . . .	5
Tibia . . . . .	3	Resection supra-orbital nerve . . . . .	2
Femur . . . . .	4	Explosion laceration face . . . . .	1
		Gangrene tongue . . . . .	1
		Hernia cerebri traumatic . . . . .	1
		Plastic mouth . . . . .	2

Noma, lip and face . . . . .	5	Dissecting cellulitis leg . . . . .	2
Compound fracture skull . . . . .	2	Perforating ulcer heel, plastic . . . . .	1
Wen of scalp . . . . .	1	Compound fracture foot . . . . .	1
Harelip . . . . .	1	Laceration of foot . . . . .	1
Empyema, vesection ribs . . . . .	3	Fistula in ano . . . . .	3
Thoracotomy . . . . .	7	Hemorrhoids . . . . .	10
Extensive specific ulcerations . . . . .	31	Circumcision . . . . .	7
Old dislocation shoulder . . . . .	2	Fissure in ano . . . . .	1
Deep infection hand . . . . .	5	Fistula and repair of rectum . . . . .	1
Sciatic foramen injection . . . . .	2	Fistula and hemorrhoids . . . . .	1
Jaw closure, arthritis . . . . .	1	Peri-rectal abscess . . . . .	2
Abscess neck and chest . . . . .	9		
Schede's operation Thoracoplasty . . . . .	1	Total miscellaneous . . . . .	151
Multiple abscesses . . . . .	1		
Mammary abscesses . . . . .	6	Summary . . . . .	
Retention cyst umbilicus . . . . .	1	Abdominal . . . . .	91
Hepatic abscess . . . . .	1	Hernias . . . . .	52
Transplantation bone splint, spine . . . . .	1	Gynecological . . . . .	65
Suppurative arthritis, shoulder . . . . .	1	Eye, ear, nose, and throat . . . . .	169
Extensive skin grafting . . . . .	1	Calculi . . . . .	10
Lacerated wound abdominal wall . . . . .	1	Amputations . . . . .	24
Fracture acromial process, compound . . . . .	1	Gun shot . . . . .	28
Caries tibia . . . . .	2	Genito-urinary . . . . .	22
Osteomyelitis tibia . . . . .	3	Tubercular bones and joints . . . . .	64
Osteomyelitis fibula . . . . .	1	Tubercular sinuses and abscesses . . . . .	66
Deep infection forearm . . . . .	1	Tumors . . . . .	45
Compound fracture tibia and fibula . . . . .	1	Miscellaneous . . . . .	151
Simple fracture tibia and fibula . . . . .	1		
Compound fracture neck humerus . . . . .	1	Total operations . . . . .	787
Caries metacarpal bones . . . . .	5		
Evulsion thigh (machinery) . . . . .	1		
Stab wound buttocks . . . . .	2		
Compound fracture thigh . . . . .	1		
Resection head of femur . . . . .	1		
Exostosis metatarsal bones . . . . .	1		
Ulceration dorsum foot . . . . .	1		
Compound fracture ankle . . . . .	1		
Double club foot, tenotomy . . . . .	2		

There have been eleven deaths in the hospital, two from Empyema, and one each from the following: Volvulus intestine of 5 days duration, Septicemia, Explosion of powder, Urinary fistula, Gastro-enterostomy followed by pulmonary oedema, Stab-wound of abdomen, Strangulated hernia gangrenous, Suprapubic cystotomy, and Pyosalpinx from shock.

PA5-14